Confidence Intervals II

Grinnell College

October 10, 2025

Warm-up

- What is the expression/formula for a confidence interval?
- ▶ What factors can change the length of my interval? How do they change it?
- ► What are some things I should consider when deciding how big to make my interval?

Review

The Law of Large Numbers guarantees that, as the number of observations n in my sample increases, my estimate of the parameter will converge to the true value

A **sampling distribution** refers to the distribution of a sample statistic (i.e., \overline{x}) if we were to repeatedly sample from a population and recompute the statistic


- What values would they take?
- ► How frequently would they appear?

The **Central Limit Theorem** states that is my statistic is an average or a proportion, then the sampling distribution of my statistic will be approximately normal, with

$$\overline{X} \sim N\left(\mu, \frac{\sigma}{\sqrt{n}}\right)$$


Grinnell College





colour —
$$\sigma = 30$$
 — $\sigma = 15$ — $\sigma = 5$

Empirical Rule



Point \pm Margin of Error

The key idea is this:

In finding a plausible range for our parameter μ , we want to create an interval, centered at our observed statistic, that takes the form

$$\overline{X} \pm C \times \left(\frac{\hat{\sigma}}{\sqrt{n}}\right)$$

Our **confidence**, mediated by C, is the frequency with which an interval constructed in this way will contain the true value of the mean.

Critical Values and Percentiles


The *C* term in the expression


$$\overline{X} \pm C \times \left(\frac{\hat{\sigma}}{\sqrt{n}}\right)$$


is called a **critical value**. The choice of *C* is the only thing we choose that determines the confidence level of our interval


Put another way: there are several things that determine the *length* of an interval, but only one that determines its confidence


Grinnell College STA 209 is cool :) October 10, 2025 7 / 20















Confidence Intervals

It is also worth observing that we can *alter* our process to achieve different results. There is a trade-off between how frequently we are correct and how much uncertainty we allow in our prediction













High Variance, C = 2









High Variance, C = 0.5







Examples

Suppose I sample n=20 male Adelie penguins from a population where $\mu=40.39$ and $\sigma=2.2771$. From my sample, I find the statistics,

$$\bar{x} = 40.63, \quad \hat{\sigma} = 1.97, \quad \frac{\hat{\sigma}}{\sqrt{n}} = 0.44$$

Find confidence intervals associated with the critical values for:

- C = 0.5
- ightharpoonup C = 1
- C = 3

Review

- ▶ Standard deviation (σ) is an estimate of the amount of variability in our sample, while standard error (σ/\sqrt{n}) is an estimate of the variability in estimating a parameter
- A sampling distribution describes the distribution of a statistic or parameter estimate if we could repeat the sampling process as many times as we wish
- Approximations to the normal distribution generally follow the **66-95-99 rule** with 1/2/3 standard deviations of the mean
- If these properties hold, we can create a reasonable interval of possible parameter values of the form Point Estimate \pm Margin of Error
- A confidence interval is an interval with the properties that:
 - It is constructed according to a procedure or set of rules
 - ▶ It is intended to give plausible range of values for a *parameter* based on a *statistic*
 - It has no probability; the interval either contains the true value or it does not

 Grinnell College
 STA 209 is cool :)
 October 10, 2025
 20 / 20